

EYE CHART

Presbyopia Patient Directions

View the eye chart without reading glasses.
Take product as directed by your eye doctor. View the eye chart again.



Talk to your eye doctor
for more information.

Font size^{1,2} Vision Test

16 pt	Presbyopia is a natural part of aging.
14 pt	You're likely not the only person reading
12 pt	an eye chart like this right now. The first
9.5 pt	symptoms most people notice are difficulty
7.5 pt	reading fine print, especially in low light conditions.
6 pt	Difficulty seeing and focusing on objects that are close to you
4.5 pt	is a common symptom. As we age, the lenses of our eyes become less flexible.

USE

VUITY® (pilocarpine hydrochloride ophthalmic solution) 1.25% is a prescription eye drop used to treat age-related blurry near vision (presbyopia) in adults.

IMPORTANT SAFETY INFORMATION

- Do not use VUITY if you are allergic to any of the ingredients.
- VUITY may cause temporary dim or dark vision. Use caution when driving at night or performing hazardous activities in poor lighting.
- Temporary problems when changing focus between near and distant objects may occur. Do not drive or use machinery if vision is not clear (e.g., blurred vision).
- Seek immediate medical care if you experience sudden onset of flashing lights, floaters, or vision loss.

Please see additional Important Safety Information on the previous page and the full Prescribing Information.



DROP BY TO SEE MORE
at Vuity.com

IMPORTANT SAFETY INFORMATION (CONTINUED)

- If you wear contact lenses, they should be removed prior to VUITY use. Wait 10 minutes after dosing before reinserting contact lenses.
- Do not touch the dropper tip to any surface as this may contaminate the contents.
- If more than one topical eye medication is being used, the medicines must be administered at least 5 minutes apart.
- The most common side effects are headache and eye redness. These are not all the possible side effects of VUITY.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

Please see Use and additional Important Safety Information on the previous page and the full [Prescribing Information](#).

Reference: 1. Sanders DR, Sanders ML. J Refract Surg. 2007;23:747-751. 2. McDonald MB, Barnett M. Ophthalmol Ther. 2022;11(1):1-11.



© 2022 AbbVie. All rights reserved.
All trademarks are the property of their respective owners.
US-VUI-220403